

Hydrogen Epoch of Reionization Array



Adrian Liu, Hubble Fellow, UC Berkeley

What is HERA?

A **funded (!)** interferometer array of 350
dishes of 14 m diameter to do low-
frequency 21 cm cosmology



154 m


What is HERA?


- 350 dishes, 14m diameter each.
- Primary science focus: $6 < z < 13$, with possibility of $5.5 < z < 27$.
- Drift-scan instrument.
- High significance measurement of 21cm power spectrum: $\sim 20\sigma$ to $\sim 90\sigma$ depending on foregrounds


HERA is primarily
designed to access the
Epoch of Reionization...


A three-parameter reionization model





- ζ : ionizing efficiency of first galaxies
- T_{vir} : minimum virial temperature (proxy for mass) of first ionizing galaxies
- R_{mfp} : mean free path of ionizing photons




























HERA is primarily designed
to access the Epoch of
Reionization...

HERA is primarily designed
to access the Epoch of
Reionization...but already
has some sensitivity to
cosmology...

Cosmological parameter uncertainties are non-negligible for HERA




Cosmo params fixed
Cosmo params varied



$$\tau \propto \int \langle (1 - x_{\text{HI}}) \delta_b \rangle dz$$


21cm information breaks the degeneracy between the amplitude of fluctuations and the optical depth



AL et al. (2016)

$$\Delta \ln(10^{10} A_s) = \pm 0.023 \longrightarrow \pm 0.0053$$

Both A_s and the neutrino mass can affect small scale power, leading to degeneracies



Allison et al. (2015)

AL et al. (2016)


$$\sum m_\nu = 60 \pm 19 \text{ meV} \longrightarrow \pm 12 \text{ meV}$$

What drives the design of HERA?
What are the similarities and differences when compared to a cosmology-centric 21cm instrument?

Why is HERA such a
regular array?

A regular grid enables:

- Calibration via redundancy
- High sensitivity by repeated measurements of the same Fourier modes




A regular grid enables:

- Calibration via redundancy
- High sensitivity by repeated measurements of the same Fourier modes

With the option of:

- Using FFT correlation




A regular grid enables:

- Calibration via redundancy
- High sensitivity by repeated measurements of the same Fourier modes

With the option of:

- Using FFT correlation





Why does HERA have
such short baselines?

- Short baselines are less chromatic
- A compact array gives better thermal noise sensitivity
- High k modes are accessed along the line-of-sight anyway

21cm interferometers are not naturally C_ℓ instruments at high z


Max. baseline length


$$\ell \sim \frac{2\pi b}{\lambda} = \frac{2\pi b \nu_{21}}{c(1+z)}$$



Why does HERA use
such big dishes?

HERA's strategy is to work at high k to avoid foregrounds. Thermal noise is high there, so large collecting areas are necessary




Why not have smaller
dishes, but have more of
them?

- Having more dishes would require a much larger correlator to get to the same collecting area.
- Larger dishes have smaller fields of view, reducing the response to the horizon.

- Having more dishes would require a much larger correlator to get to the same collecting area.
- Larger dishes have smaller fields of view, reducing the response to the horizon.
- This is not without downsides:
 - Cosmic variance? Likely not a problem.
 - Limited cross-correlation options with other surveys.

HERA is optimized for its science case,
but various design parameters should be
revisited for futuristic arrays



- Larger field-of-view?
- Tracking?
- Faster (e.g. FFT) correlators?
- Will longer baselines/angular information become more necessary?